

COMMUNICATIONS

Mathematics

SPLITTING AUTOMORPHISMS OF FREE BURNSIDE GROUPS

V. S. ATABEKYAN*

Chair of Algebra and Geometry, YSU

We have proved, that if $n \geq 1003$ is an arbitrary odd number and φ is a splitting automorphism of period n of group $B(m, n)$ that has a prime order, then φ is inner automorphism.

Keywords: splitting automorphism, free Burside group, holomorph of the group.

Definition. An automorphism φ of G is called *splitting automorphism of period n* , if $\varphi^n = 1$ and $g \varphi(g) \varphi^2(g) \dots \varphi^{n-1}(g) = 1$ for each $g \in G$.

If φ is a splitting automorphism of period n of a group G , then for every $g \in G$ the relation $(\varphi g)^n = 1$ holds in the holomorph $Hol(G)$ of the group G . In particular, the identity automorphism of G is a splitting automorphism of period n , iff the identity $x^n = 1$ is satisfied in G .

A well-known Theorem of O. Kegel states that any finite group that has a nontrivial splitting automorphism of prime period is nilpotent (see [1]). This result generalizes J. Tompson's Theorem [2] on the nilpotency of finite group with an automorphism of prime order without fixed points. E. Khukhro proved that any solvable group having a nontrivial splitting automorphism of prime period is also a nilpotent group. (see [3]).

If Γ is any group satisfying the identical relation $x^n = 1$, then the identities

$$b(a^{-1}ba)(a^{-2}ba^2) \dots (a^{-n+1}ba^{n-1})^n a^n, \quad a^{-n}ba^n = b$$

also hold in the group Γ . Consequently, each inner automorphism $i_a \in \text{Inn}(\Gamma)$, $i_a(b) = a^{-1}ba$ is a splitting automorphism of period n of the group Γ .

We are interested in the inverse question for splitting automorphisms of the free Burnside group $B(m, n)$. By definition, a free Burnside group $B(m, n)$ of period n and rank m has the following presentation

$$B(m, n) = \langle a_1, a_2, \dots, a_m \mid X^n = 1 \rangle,$$

where X runs over all words in $\{a_1^{\pm 1}, a_2^{\pm 1}, \dots, a_m^{\pm 1}\}$.

* E-mail: varujan@atabekyan.com

The group $B(m, n)$ is the quotient of the free group F_m of rank m by the normal subgroup F_m^n , generated by all n th powers of elements of F_m . Each periodic group of period n with m generators is a quotient group of $B(m, n)$. According to the fundamental Theorem of S.I. Adian that solves the Burnside problem (see [4]), for any odd $n \geq 665$ and $m > 1$ the group $B(m, n)$ is infinite. A detailed review of studies on free Burnside groups is given by S.I. Adian in [5].

In 1991 S.V. Ivanov posed the following question: let $m > 1$ and n is large enough odd number. Is it true that any splitting automorphism φ of $B(m, n)$ is inner (see Kourovka notebook [6], question 11.36, b)?

We have proved the following theorem.

Theorem. Let $n \geq 1003$ be an arbitrary odd number and φ be a splitting automorphism of period n of $B(m, n)$. If the order of the automorphism φ is a prime number, then φ is inner.

This Theorem immediately implies.

Corollary. For any prime $n > 997$ and $m > 1$ each splitting automorphism of the group $B(m, n)$ is an inner automorphism.

Outline of the Proof of the Main Result. In the paper [7] it was proved that for any $m > 1$ and odd $n \geq 1003$ there exists a maximal normal subgroup N of the free Burnside group $B(m, n)$, such that the quotient $B(m, n)/N$ is an infinite group, every proper subgroup of which is contained in some cyclic subgroup of order n . Denote by \mathcal{M}_n the set of all such maximal normal subgroups N of the free Burnside group $B(m, n)$. The groups $B(m, n)/N$, constructed in [7], are called Tarski monsters. In [8] it was shown that for every odd $n \geq 1003$ there is a continuum of non-isomorphic Tarski monsters of period n .

The following statement, proved by the author in [9], plays a key role in the proof of the main result.

Proposition 1. (see [9], Corollary 2) Let $n \geq 1003$ be an arbitrary odd number and φ be an automorphism of the group $B(m, n)$, such that $\varphi(N) = N$ for any $N \in \mathcal{M}_n$. Then φ is inner automorphism.

The following interesting results obtained by the author are also used in the proof.

Proposition 2. Let $\varphi: G \rightarrow G$ be an arbitrary automorphism and N be a normal subgroup of G , such that the quotient G/N is a non-abelian simple group. If the subgroups $N, \varphi(N), \dots, \varphi^{k-1}(N)$ are pairwise distinct and $\varphi^k(N) = N$, then

the quotient group $G / \bigcap_{i=1}^k \varphi^i(N)$ is decomposed into a direct product of subgroups

$N_j / \bigcap_{i=1}^k \varphi^i(N)$, $j = 1, 2, \dots, k$, where $N_j = \bigcap_{i=1, i \neq j}^k \varphi^i(N)$. Moreover the quotient

$N_j / \bigcap_{i=1}^k \varphi^i(N)$ is isomorphic to G/N .

Proposition 3. If $n \geq 1003$ is an odd number and φ is an arbitrary nontrivial splitting automorphism of period n of $B(m, n)$, then for every normal subgroup $N \in \mathcal{M}_n$ the stabilizer relative to action of the cyclic group $\langle \varphi \rangle$ is nontrivial.

Received 14.04.2011

REFERENCES

1. **Kegel O.H.** Math. Z., 1961, v. 75, p. 373–376.
2. **Thompson J.G.** Proc. Nat. Acad. Sci. USA, 1959, v. 45, p. 578–581.
3. **Khukhro E.I.** Algebra i Logika, 1980, v. 19, № 1, p. 118–129.
4. **Adian S.I.** The Burnside Problem and Identities in Groups. *Ergeb. Math. Grenzgeb. V. 95.* Berlin–New York: Springer–Verlag, 1979.
5. **Adyan S.I.** Russian Math. Surveys, 2010, v. 65, № 5, p. 805–855.
6. **Mazurov V.D., Merzlyakov Yu.I., Churkin V.A.** (eds.). The Kourovka Notebook. Unsolved Problems in Group Theory (ed. 11). Institute of Math. Novosibirsk, 1990.
7. **Adian S.I., Lysenok I.G.** Math. USSR. Izv., 1992, v. 39, № 2, p. 905–957.
8. **Atabekyan V.S.** Mathematical Notes, 2007, v. 82, № 4, p. 443–447.
9. **Atabekyan V.S.** Izv. RAN. Ser. Mat., 2011, v. 75, № 2, p. 3–18 (in Russian)

Վ. Ս. Աթաբեկյան. Ազատ բեռնասայրյան խմբի տրոհող ավտոմորֆիզմներ էջ. 62–64

Ապացուցվել է, որ եթե $n \geq 1003$ կենս թիվ է և φ -ն $B(m, n)$ խմբի պարբերությամբ տրոհող ավտոմորֆիզմ, որն ունի պարզ կարգ, ապա φ -ն ներքին ավտոմորֆիզմ է:

В. С. Атабекян. Расщепляющие автоморфизмы свободных бернсайдовых групп
стр. 62–64

Доказано, что если $n \geq 1003$ – произвольное нечетное число и φ – произвольный расщепляющий автоморфизм периода n группы $B(m, n)$, который имеет простой порядок, то φ является внутренним автоморфизмом.