

Mathematics

ON BOUNDED OPERATORS IN  $L^p$  SPACES

A. I. PETROSYAN\*, A. F. BEKNAZARYAN

Chair of Mathematical Analysis, YSU

In the present paper the linear operators that depend on a normal pair of weight functions  $\{\varphi, \psi\}$  in the Banach spaces  $L^p(D)$ , where  $D$  is the unit disk in the complex plane, are considered. It is investigated, for which values of  $p$  these operators are bounded.

**Keywords:** Banach space, analytic function, bounded operator.

**Introduction.** The paper is devoted to investigation of some linear operators depending on a normal pair of weight functions in  $L^p(D)$  spaces. The concept of a normal pair of weight functions introduced for the first time by Shields and Williams [1] and was appeared convenient for statement of the estimates of integrals and for exposition of projectors in weight spaces. The basic result of paper is formulated in Theorem 2. The case  $p=1$  is discussed in [1]. Note that the case of power weight functions is considered in [2] (Chapter 1, § 2, Theorem 1.9).

We use the following notations:  $D = \{z \in \mathbb{C} : |z| < 1\}$  is the unit disk in the complex plane  $\mathbb{C}$ ,  $dA$  is the normalized area measure on  $D$ . In terms of real coordinates we have  $dA(z) = \frac{1}{\pi} dx dy = \frac{1}{\pi} r dr d\theta$ ,  $z = x + iy = re^{i\theta}$ .

We define  $L^p(D, dA)$  as the set of all functions  $f$  that are measurable by  $dA$  in  $D$ , for which

$$\|f\|_p = \left\{ \int_D |f(w)|^p dA(w) \right\}^{1/p} < +\infty, \quad 0 < p < \infty.$$

Let the functions  $\varphi(r)$  and  $\psi(r)$  be positive and continuous on  $(0,1]$  with

$$\lim_{r \rightarrow 1} \varphi(r) = 0 \quad \text{and} \quad \int_0^1 \psi(r) dr < \infty.$$

**Definition 1.** A function  $\varphi(r)$  is called normal, if there exist  $k > \varepsilon > 0$  and  $r_0 < 1$  such that

$$\frac{\varphi(r)}{(1-r)^\varepsilon} \searrow 0, \quad \frac{\varphi(r)}{(1-r)^k} \nearrow \infty \quad (r_0 \leq r, \quad r \rightarrow 1-0). \quad (1)$$

\* E-mail: [apetrosyan@ysu.am](mailto:apetrosyan@ysu.am)

**Definition 2.** The functions  $\{\varphi, \psi\}$  is called a normal pair, if  $\varphi$  is normal and if for some  $k$  satisfying (1), there exists  $\alpha > k - 1$  such that

$$\varphi(r)\psi(r) = (1-r^2)^\alpha, \quad 0 \leq r < 1.$$

Consider the integral operator

$$Tf(z) = \int_D \frac{\psi(z)\varphi(w)}{|1-z\bar{w}|^{2+\alpha}} f(w) dA(w). \quad (2)$$

We want to establish for which  $p$  this operator is bounded.

**Auxiliary Lemmas.**

**Lemma 1.** For  $\gamma > -1$  and  $m > 1 + \gamma$  we have

$$\int_0^1 (1-\rho r)^{-m} (1-r)^\gamma dr \leq C(1-\rho)^{1+\gamma-m}, \quad 0 < \rho < 1,$$

where the constant  $C = C(m, \gamma)$  does not depend on  $\rho$ . For the proof see [1] (Chapter 1, § 3).

**Lemma 2.** For a positive  $\beta > 0$  and  $z \in D$

$$\int_0^{2\pi} \frac{d\theta}{|1-ze^{-i\theta}|^{1+\beta}} = O\left(\frac{1}{(1-|z|^2)^\beta}\right).$$

For the proof see [3] (Chapter 4, § 6).

**Lemma 3.** For  $k - \alpha < \frac{\beta}{p} < 1 + \varepsilon$  there exist a constant  $M_0$  such that

$$\int_D \frac{\varphi(w)}{(1-|w|^2)^{\beta/p} |1-z\bar{w}|^{2+\alpha}} dA(w) \leq M_0 \frac{1}{\psi(z)(1-|z|^2)^{\beta/p}}.$$

*Proof.* According to the definition of a normal pair of functions,  $\varphi(r)\psi(r) = (1-r^2)^\alpha$  for  $0 \leq r < 1$ , and it will be sufficient to show that

$$\int_D \frac{\varphi(w)}{(1-|w|^2)^{\beta/p} |1-z\bar{w}|^{2+\alpha}} dA(w) \leq M_0 \frac{\varphi(z)}{(1-|z|^2)^{(\beta/p)+\alpha}}.$$

Let  $w = re^{i\theta}$ . As  $\alpha + 1 > 0$ , then, due to Lemma 2, there exists a constant  $C$  such that  $\int_0^{2\pi} \frac{1}{|1-zre^{-i\theta}|^{\alpha+2}} d\theta \leq \frac{C}{(1-(|z|r)^2)^{\alpha+1}}$ .

Passing to polar coordinates, we receive

$$\begin{aligned} \int_D \frac{\varphi(w)}{(1-|w|^2)^{\beta/p} |1-z\bar{w}|^{2+\alpha}} dA(w) &= \frac{1}{\pi} \int_0^1 \frac{r\varphi(r)}{(1-r^2)^{\beta/p}} \int_0^{2\pi} \frac{d\theta}{|1-zre^{-i\theta}|^{\alpha+2}} dr \leq \\ &\leq C \int_0^1 \frac{\varphi(r)}{(1-|r|^2)^{\beta/p} (1-(|z|r)^2)^{\alpha+1}} dr \leq C_1 \int_0^1 \frac{\varphi(r)}{(1-|r|)^{\beta/p} (1-|z|r)^{\alpha+1}} dr. \end{aligned} \quad (3)$$

Divide the last integral into three parts:

$$\begin{aligned} \int_0^1 \frac{\varphi(r)}{(1-r)^{\beta/p} (1-|z|r)^{\alpha+1}} dr &= \int_0^{r_0} \frac{\varphi(r)}{(1-r)^{\beta/p} (1-|z|r)^{\alpha+1}} dr + \\ &+ \int_{r_0}^{|z|} \frac{\varphi(r)}{(1-r)^{\beta/p} (1-|z|r)^{\alpha+1}} dr + \int_{|z|}^1 \frac{\varphi(r)}{(1-r)^{\beta/p} (1-|z|r)^{\alpha+1}} dr = I_1 + I_2 + I_3. \end{aligned} \quad (4)$$

Obviously  $I_1$  is bounded for all  $z$ . Therefore, there exists a constant  $C_2$  such that

$$I_1 \leq C_2 \frac{\varphi(z)}{(1-|z|^2)^{(\beta/p)+\alpha}}. \tag{5}$$

From the definition of normal function (1) we have  $\frac{\varphi(r)}{(1-r)^k} \leq \frac{\varphi(z)}{(1-|z|)^k}$  for

$r_0 < r \leq |z|$ . Therefore,

$$I_2 = \int_{r_0}^{|z|} \frac{\varphi(r)}{(1-r)^{\beta/p} (1-|z|r)^{\alpha+1}} dr = \int_{r_0}^{|z|} \frac{\varphi(r)(1-r)^{k-(\beta/p)}}{(1-r)^k (1-|z|r)^{\alpha+1}} dr \leq \frac{\varphi(z)}{(1-|z|)^k} \int_{r_0}^{|z|} \frac{(1-r)^{k-(\beta/p)}}{(1-|z|r)^{\alpha+1}} dr.$$

As  $k - \alpha < \frac{\beta}{p} < 1 + \varepsilon$ , then  $k - \frac{\beta}{p} < \alpha$  and  $-1 < \varepsilon - \frac{\beta}{p} < k - \frac{\beta}{p}$ , that is

$-1 < k - \frac{\beta}{p} < \alpha$ , and taking into account Lemma 1 we have

$$\int_{r_0}^{|z|} \frac{(1-r)^{k-(\beta/p)}}{(1-|z|r)^{\alpha+1}} dr \leq \int_0^1 \frac{(1-r)^{k-(\beta/p)}}{(1-|z|r)^{\alpha+1}} dr \leq C_3 \frac{1}{(1-|z|)^{(\beta/p)+\alpha-k}}$$

for some constant  $C_3$ . Therefore,

$$I_2 \leq \frac{\varphi(z)}{(1-|z|)^k} \int_{r_0}^{|z|} \frac{(1-r)^{k-(\beta/p)}}{(1-|z|r)^{\alpha+1}} dr \leq \frac{\varphi(z)}{(1-|z|)^k} \frac{C_3}{(1-|z|)^{(\beta/p)+\alpha-k}} \leq C_4 \frac{\varphi(z)}{(1-|z|^2)^{(\beta/p)+\alpha}}. \tag{6}$$

Now we shall pass to  $I_3$ . As  $\frac{\varphi(r)}{(1-r)^\varepsilon} \leq \frac{\varphi(z)}{(1-|z|)^\varepsilon}$  for  $|z| < r < 1$ , then

$$\begin{aligned} I_3 &= \int_{|z|}^1 \frac{\varphi(r)}{(1-r)^{(\beta/p)} (1-|z|r)^{\alpha+1}} dr = \int_{|z|}^1 \frac{\varphi(r)(1-r)^{\varepsilon-(\beta/p)}}{(1-r)^\varepsilon (1-|z|r)^{\alpha+1}} dr \leq \\ &\leq \frac{\varphi(z)}{(1-|z|)^\varepsilon} \int_{|z|}^1 \frac{(1-r)^{\varepsilon-(\beta/p)}}{(1-|z|r)^{\alpha+1}} dr \leq \frac{\varphi(z)}{(1-|z|)^\varepsilon} \int_0^1 \frac{(1-r)^{\varepsilon-(\beta/p)}}{(1-|z|r)^{\alpha+1}} dr. \end{aligned}$$

We have also  $k - \alpha < \frac{\beta}{p} < 1 + \varepsilon$ , therefore,  $\varepsilon - \frac{\beta}{p} > -1$  and  $\varepsilon - \alpha < k - \alpha < \frac{\beta}{p}$ , i.e.

$\alpha + 1 > 1 + \varepsilon - \frac{\beta}{p}$ . Again using Lemma 1 we get

$$\int_0^1 \frac{(1-r)^{\varepsilon-(\beta/p)}}{(1-|z|r)^{\alpha+1}} dr \leq C_5 \frac{1}{(1-|z|)^{(\beta/p)+\alpha-\varepsilon}},$$

whence we have

$$I_3 \leq \frac{\varphi(z)}{(1-|z|)^\varepsilon} \int_0^1 \frac{(1-r)^{\varepsilon-(\beta/p)}}{(1-|z|r)^{\alpha+1}} dr \leq C_5 \frac{\varphi(z)}{(1-|z|)^{(\beta/p)+\alpha}} \leq C_6 \frac{\varphi(z)}{(1-|z|^2)^{(\beta/p)+\alpha}} \tag{7}$$

with some constants  $C_5$  and  $C_6$ . Combining the obtained results (3)–(7), we get the constant  $M_0$  such that

$$\int_D \frac{\varphi(w)}{(1-|w|^2)^{(\beta/p)} |1-z\bar{w}|^{2+\alpha}} dA(w) \leq M_0 \frac{\varphi(z)}{(1-|z|^2)^{(\beta/p)+\alpha}} = M_0 \frac{1}{\psi(z)(1-|z|^2)^{\beta/p}}$$

for  $k - \alpha < \frac{\beta}{p} < 1 + \varepsilon$ .

The Lemma is proved.

**Lemma 4.** Let  $-\varepsilon < \frac{\beta}{q} < \alpha + 1 - k$ . There exists a constant  $M_1$  such that

$$\int_D \frac{\psi(z)}{(1-|z|^2)^{\beta/p} |1-z\bar{w}|^{2+\alpha}} dA(z) \leq M_1 \frac{1}{\varphi(w)(1-|w|^2)^{\beta/p}}.$$

*Proof.* For  $0 \leq r < 1$  we have  $\varphi(r)\psi(r) = (1-r^2)^\alpha$ . Therefore, it is necessary to prove that

$$\int_D \frac{(1-|z|^2)^{\alpha-(\beta/p)}}{\varphi(z)|1-z\bar{w}|^{2+\alpha}} dA(z) \leq M_1 \frac{1}{\varphi(w)(1-|w|^2)^{\beta/p}}$$

for any constant  $M_1$ . Let  $z = re^{i\theta}$ . Then

$$\int_D \frac{(1-|z|^2)^{\alpha-(\beta/p)}}{\varphi(z)|1-z\bar{w}|^{2+\alpha}} dA(z) = \frac{1}{\pi} \int_0^1 \frac{r(1-r^2)^{\alpha-(\beta/p)}}{\varphi(r)} \int_0^{2\pi} \frac{d\theta}{|1-\bar{w}re^{i\theta}|^{\alpha+2}} dr,$$

and, due to Lemma 2, there exists a constant  $K_1$  such that

$$\begin{aligned} \frac{1}{\pi} \int_0^1 \frac{r(1-r^2)^{\alpha-(\beta/p)}}{\varphi(r)} \int_0^{2\pi} \frac{d\theta}{|1-\bar{w}re^{i\theta}|^{\alpha+2}} dr &= \frac{1}{\pi} \int_0^1 \frac{r(1-r^2)^{\alpha-(\beta/p)}}{\varphi(r)} \int_0^{2\pi} \frac{d\theta}{|1-wre^{-i\theta}|^{\alpha+2}} dr \leq \\ &\leq K_1 \int_0^1 \frac{(1-r^2)^{\alpha-(\beta/p)}}{\varphi(r)(1-(|w|r)^2)^{\alpha+1}} dr \leq K_2 \int_0^1 \frac{(1-r)^{\alpha-(\beta/p)}}{\varphi(r)(1-|w|r)^{\alpha+1}} dr \end{aligned}$$

(we have replaced  $(1-r^2)$  by  $(1-r)$  and  $(1-(|w|r)^2)$  by  $(1-|w|r)$ , therefore, the value of  $K_1$  was replaced by some other value of  $K_2$ ). Dividing the last integral into three parts, we obtain

$$\begin{aligned} \int_0^1 \frac{(1-r)^{\alpha-(\beta/p)}}{\varphi(r)(1-|w|r)^{\alpha+1}} dr &= \int_0^{r_0} \frac{(1-r)^{\alpha-(\beta/p)}}{\varphi(r)(1-|w|r)^{\alpha+1}} dr + \\ &+ \int_{r_0}^{|w|} \frac{(1-r)^{\alpha-(\beta/p)}}{\varphi(r)(1-|w|r)^{\alpha+1}} dr + \int_{|w|}^1 \frac{(1-r)^{\alpha-(\beta/p)}}{\varphi(r)(1-|w|r)^{\alpha+1}} dr = J_1 + J_2 + J_3. \end{aligned}$$

Integral  $J_1$  is bounded for all  $w$ , therefore, there is  $K_3$  such that

$$J_1 \leq \frac{K_3}{\varphi(w)(1-|w|^2)^{\beta/p}}. \quad (8)$$

From the definition of normal function (1) we have  $\frac{(1-r)^\varepsilon}{\varphi(r)} \leq \frac{(1-|w|)^\varepsilon}{\varphi(w)}$ , if

$r_0 < r \leq |w|$ . Therefore,

$$\begin{aligned} J_2 &= \int_{r_0}^{|w|} \frac{(1-r)^{\alpha-(\beta/p)}}{\varphi(r)(1-|w|r)^{\alpha+1}} dr = \int_r^{|w|} \frac{(1-r)^\varepsilon (1-r)^{\alpha-(\beta/p)-\varepsilon}}{\varphi(r)(1-|w|r)^{\alpha+1}} dr \leq \\ &\leq \frac{(1-|w|)^\varepsilon}{\varphi(w)} \int_{r_0}^{|w|} \frac{(1-r)^{\alpha-(\beta/p)-\varepsilon}}{(1-|w|r)^{\alpha+1}} dr \leq \frac{(1-|w|)^\varepsilon}{\varphi(w)} \int_0^1 \frac{(1-r)^{\alpha-(\beta/p)-\varepsilon}}{(1-|w|r)^{\alpha+1}} dr. \end{aligned}$$

As  $-\varepsilon < \frac{\beta}{q} < \alpha + 1 - k$ , then  $\alpha - \frac{\beta}{q} - \varepsilon > \alpha - \frac{\beta}{q} - k > -1$  and  $\alpha + 1 > \alpha - \frac{\beta}{q} - \varepsilon + 1$ .

Using Lemma 1 we obtain  $\int_0^1 \frac{(1-r)^{\alpha-(\beta/p)-\varepsilon}}{(1-|w|r)^{\alpha+1}} dr \leq K_4 \frac{1}{(1-|w|)^{\varepsilon+(\beta/p)}}$  and, therefore,

$$J_2 \leq \frac{(1-|w|)^\varepsilon}{\varphi(w)} \cdot \frac{K_4}{(1-|w|)^{\varepsilon+(\beta/p)}} = \frac{K_4}{\varphi(w)(1-|w|)^{\beta/p}} \leq \frac{K_5}{\varphi(w)(1-|w|^2)^{\beta/p}}. \quad (9)$$

For estimation of  $J_3$  we use the inequality  $\frac{(1-r)^k}{\varphi(r)} \leq \frac{(1-|w|)^k}{\varphi(w)}$ , if  $|w| < r < 1$ .

$$\begin{aligned} J_3 &= \int_{|w|}^1 \frac{(1-r)^{\alpha-(\beta/p)}}{\varphi(r)(1-|w|r)^{\alpha+1}} dr = \int_{|w|}^1 \frac{(1-r)^k (1-r)^{\alpha-(\beta/p)-k}}{\varphi(r)(1-|w|r)^{\alpha+1}} dr \leq \\ &\leq \frac{(1-|w|)^k}{\varphi(w)} \int_{|w|}^1 \frac{(1-r)^{\alpha-(\beta/p)-k}}{(1-|w|r)^{\alpha+1}} dr \leq \frac{(1-|w|)^k}{\varphi(w)} \int_0^1 \frac{(1-r)^{\alpha-(\beta/p)-k}}{(1-|w|r)^{\alpha+1}} dr. \end{aligned}$$

From  $-\varepsilon < \frac{\beta}{q} < \alpha + 1 - k$  we have  $\alpha - k - \frac{\beta}{q} > -1$  and  $\alpha - k - \frac{\beta}{q} + 1 < \alpha - \varepsilon - \frac{\beta}{q} + 1 < \alpha + 1$ .

Taking into account Lemma 1 we get  $\int_0^1 \frac{(1-r)^{\alpha-(\beta/p)-k}}{(1-|w|r)^{\alpha+1}} dr \leq K_6 \frac{1}{(1-|w|)^{k+(\beta/p)}}$  for a constant  $K_6$ . Therefore,

$$J_3 \leq \frac{(1-|w|)^k}{\varphi(w)} \cdot \frac{K_6}{(1-|w|)^{k+(\beta/p)}} = \frac{K_6}{\varphi(w)(1-|w|)^{\beta/p}} \leq \frac{K_7}{\varphi(w)(1-|w|^2)^{\beta/p}}. \quad (10)$$

Combining the obtained inequalities (7)–(10) for  $J_1$ ,  $J_2$  and  $J_3$ , we get the constant  $M_1$  such that  $\int_D \frac{\psi(z)}{(1-|z|^2)^{\beta/p} |1-z\bar{w}|^{2+\alpha}} dA(z) \leq M_1 \frac{1}{\varphi(w)(1-|w|^2)^{\beta/p}}$ .

The Lemma is proved.

**The Main Result.** The following Schur test is a useful tool for estimation of  $L^p$ .

**Theorem 1** (Schur test). Suppose  $(X, \mu)$  is a measure space,  $1 < p < \infty$ , and  $\frac{1}{p} + \frac{1}{q} = 1$ . For a nonnegative kernel  $H(x, y)$  consider the integral operator

$$(Tf)(x) = \int_X H(x, y) f(y) d\mu(y).$$

If there exist a positive function  $h$  on  $X$  and a positive constant  $C$  such that

$$\int_X H(x, y) h(y)^q d\mu(y) \leq Ch(x)^q$$

for almost all  $x \in X$  and

$$\int_X H(x, y) h(x)^p d\mu(x) \leq Ch(y)^p$$

for almost all  $y \in X$ , then the operator  $T$  is bounded on  $L^p(X, \mu)$  with  $\|T\| \leq C$ .

**Theorem 2.** For  $p(k-\alpha) < 1$  the integral operator (2) is bounded in  $L^p(D, dA)$ .

*Proof.* According to the Schur test, it would be sufficient to find an approximation function  $h$ , for which the inequalities

$$\int_D \frac{\psi(z)\varphi(w)}{|1-z\bar{w}|^{2+\alpha}} h(w)^q dA(w) \leq Mh(z)^q$$

and

$$\int_D \frac{\psi(z)\varphi(w)}{|1-z\bar{w}|^{2+\alpha}} h(z)^p dA(z) \leq Mh(w)^p$$

are fulfilled. We find the function  $h$ , having  $h(z) = \frac{1}{(1-|z|^2)^{\beta/pq}}$  form. That is

we should prove that

$$\int_D \frac{\varphi(w)}{(1-|w|^2)^{\beta/p} |1-z\bar{w}|^{2+\alpha}} dA(w) \leq M \frac{1}{\psi(z)(1-|z|^2)^{\beta/p}} \quad (11)$$

and

$$\int_D \frac{\psi(z)}{(1-|z|^2)^{\beta/p} |1-z\bar{w}|^{2+\alpha}} dA(z) \leq M \frac{1}{\varphi(w)(1-|w|^2)^{\beta/p}}. \quad (12)$$

According to Lemmas 3 and 4, the inequalities (11) and (12) are valid under the conditions  $k - \alpha < \frac{\beta}{p} < 1 + \varepsilon$ ,  $-\varepsilon < \frac{\beta}{q} < \alpha + 1 - k$ . Here  $M = \max\{M_0, M_1\}$ .

Taking into account that  $q = \frac{p}{p-1}$ , we rewrite these conditions as

$$(k - \alpha)p < \beta < (1 + \varepsilon)p, \quad -\varepsilon \frac{p}{p-1} < \beta < (\alpha + 1 - k) \frac{p}{p-1}.$$

It is only necessary to establish, at what values of  $p$  the intersection of intervals

$$((k - \alpha)p, (1 + \varepsilon)p), \quad \left( -\varepsilon \frac{p}{p-1}, (\alpha + 1 - k) \frac{p}{p-1} \right)$$

is nonempty. As  $p > 1$ , hence always  $-\varepsilon \frac{p}{p-1} < (1 + \varepsilon)p$ . Therefore, it is necessary

to clarify when  $(\alpha + 1 - k) \frac{p}{p-1} > (k - \alpha)p$ . From this inequality we obtain

$p(k - \alpha) < 1$ . That is why for  $p(k - \alpha) < 1$  the operator (2) is bounded in  $L^p(D, dA)$ .

The Theorem is proved.

Received 29.12.2010

#### REFERENCES

1. **Shields A.L., Williams D.L.** Trans. Amer. Math. Soc., 1971, v. 162, p. 287–302.
2. **Hedenmalm H., Korenblum B., Zhu K.** Theory of Bergman Spaces. Springer Verlag, 2000.
3. **Duren P.L.** Theory of  $H^p$  Spaces. New York: Academic Press, 1970.

Ա. Ի. Պետրոսյան, Ա. Ֆ. Բեկնազարյան.

$L^p$  Տարածություններում սահմանափակ օպերատորների մասին

Հոդվածում հետազոտվում են  $L^p(D)$  բանախյան տարածություններում գծային օպերատորներ, որոնք կախված են կշռային ֆունկցիաների  $\{\varphi, \psi\}$  նորմալ զույգից: Այստեղ  $D$ -ն միավոր շրջանն է կոմպլեքս հարթության վրա: Պարզաբանվում է, թե որ  $p$ -երի համար այդ օպերատորները կլինեն սահմանափակ:

А. И. Петросян, А. Ф. Бекназарян.

Об ограниченных операторах в пространствах  $L^p$

В статье исследуются линейные операторы в банаховых пространствах  $L^p(D)$ , зависящие от нормальной пары весовых функций  $\{\varphi, \psi\}$ . Здесь  $D$  — единичный круг на комплексной плоскости. Выясняется, для каких  $p$  эти операторы являются ограниченными.