

Mathematics

ON COMPACTNESS OF A CLASS OF FIRST ORDER LINEAR DIFFERENTIAL OPERATORS

V. Zh. DUMANYAN*

Chair of Numerical Analysis and Mathematical Modeling, YSU

In the present article a class of first order linear differential operators with unbounded coefficients is investigated. The compactness of operators is proved.

Keywords: bounded differential operator, compact differential operator, first order differential operator.

Let $Q \subset R^n$, $n \geq 2$, be a bounded domain with smooth boundary $\partial Q \in C^1$. Consider the first order differential expression

$$Tu \equiv (\bar{b}(x), \nabla u(x)) - \operatorname{div}(\bar{c}(x)u(x)) + d(x)u(x), \quad u \in \overset{\circ}{W}_2^1(Q),$$

with coefficients $\bar{b}(x) = (b^{(1)}(x), \dots, b^{(n)}(x))$, $\bar{c}(x) = (c^{(1)}(x), \dots, c^{(n)}(x))$ and $d(x)$ that are measurable and bounded on each strong inner subdomain of the domain Q .

For an arbitrary $u, v \in \overset{\circ}{W}_2^1(Q)$ define

$$\langle Tu, v \rangle \equiv \int_Q ((\bar{b}(x), \nabla u(x))v(x) + (\bar{c}(x)u(x), \nabla v(x)) + d(x)u(x)v(x))dx, \quad v \in \overset{\circ}{W}_2^1(Q).$$

Assume that the coefficients $\bar{b}(x)$, $\bar{c}(x)$ and $d(x)$ satisfy the conditions

$$|\bar{b}(x)| = O\left(\frac{1}{r(x)}\right) \text{ as } r(x) \rightarrow 0,$$

where $r(x)$ is the distance of a point $x \in Q$ from the boundary ∂Q ,

$$\int_0^t C^2(t)dt < \infty \quad \text{with } C(t) = \sup_{r(x) \geq t} |\bar{c}(x)|,$$

$$\int_0^t D^2(t)dt < \infty \quad \text{with } D(t) = \sup_{r(x) \geq t} |d(x)|.$$

In [1] it was shown that T is a bounded linear operator from $\overset{\circ}{W}_2^1(Q)$ into $\overset{\circ}{W}_2^{-1}(Q)$. The aim of this article is to obtain conditions on coefficients $\bar{b}(x), \bar{c}(x)$

* E-mail: duman@ysu.am

and $d(x)$, for which T is a linear compact operator from $\overset{\circ}{W}_2^1(Q)$ into $\overset{\circ}{W}_2^{-1}(Q)$. This property has important applications in studying the solvability of the problems of mathematical physics, see, for example, [2, 3].

We prove the below theorem.

Theorem. Let the below conditions hold

$$|\bar{b}(x)| = o\left(\frac{1}{r(x)}\right) \text{ as } r(x) \rightarrow 0, \quad (1)$$

and there exist monotone functions $\omega_i(t) \rightarrow 0$, as $t \rightarrow +0$, $i=1,2$, such that

$$\int_0^t \frac{tC^2(t)}{\omega_1(t)} dt < \infty, \text{ where } C(t) = \sup_{r(x) \geq t} |\bar{c}(x)|, \quad (2)$$

$$\int_0^t \frac{t^3 D^2(t)}{\omega_2(t)} dt < \infty, \text{ where } D(t) = \sup_{r(x) \geq t} |d(x)|. \quad (3)$$

Then the operator T is a compact linear operator from $\overset{\circ}{W}_2^1(Q)$ into $\overset{\circ}{W}_2^{-1}(Q)$.

Proof of Theorem. We shall follow the scheme of proof of the theorem from [1].

Let $x^0 \in \partial Q$ be an arbitrary point of the boundary ∂Q of the domain Q and (x', x_n) be a local coordinate system with the origin x^0 and the x_n axis directed along the inner normal $\nu(x^0)$ to ∂Q at the point x^0 . Since $\partial Q \in C^1$, there exists a positive number $r_{x^0} > 0$ and a function $\varphi_{x^0} \in C^1(R^{n-1})$ with properties

$$\varphi_{x^0}(0) = 0, \quad \nabla \varphi_{x^0}(0) = 0 \text{ and } |\nabla \varphi_{x^0}(x')| \leq \frac{1}{2} \text{ for all } x' \in R^{n-1},$$

such that the intersection of the domain Q with the ball $U_{x^0}^{(r_{x^0})} = \{x : |x - x^0| < r_{x^0}\}$ of radius r_{x^0} and the centre x^0 has the form $Q \cap U_{x^0}^{(r_{x^0})} = U_{x^0}^{(r_{x^0})} \cap \{(x', x_n) : x_n > \varphi_{x^0}(x')\}$.

Then $\partial Q \cap U_{x^0}^{(r_{x^0})} = U_{x^0}^{(r_{x^0})} \cap \{(x', x_n) : x_n = \varphi_{x^0}(x')\}$. Let $l_{x^0} = \frac{r_{x^0}}{\sqrt{2}}$.

From the covering $\{U_{x^0}^{(l_{x^0})}, x^0 \in \partial Q\}$ of the boundary ∂Q select a finite subcovering $U_{x^m}^{(l_{x^m})}$, $m=1, \dots, p$. Denote for simplicity $U_{x^m}^{(l_{x^m})}$ by U_m , r_{x^m} by r_m , l_{x^m} by l_m , φ_{x^m} by φ_m , $m=1, \dots, p$. Set $h = \frac{1}{3} \left(\frac{2}{\sqrt{5}} - \frac{\sqrt{2}}{2} \right) \min(r_1, \dots, r_p)$. Then each of the curvilinear

“cylinders” $\prod_m^{l_m, h} = \{(x', x_n) : |x'| < l_m, \varphi_m(x') < x_n < \varphi_m(x') + h\}$, $m=1, \dots, p$,

is contained in the corresponding ball U_m , and also by $U_m \cap Q$ (recall that (x', x_n) are the coordinates of a point in a local system of coordinates with origin at x^m). Let $l_0 < h$ be such a positive number that the complement of the domain $Q_{l_0} = \{x \in Q : r(x) = \text{dist}(x, \partial Q) > l_0\}$ in Q is contained in the union of the

“cylinders” $\prod_m^{l_m, h}$, $m=1, \dots, p$, i.e. $Q^{l_0} = \{x \in Q : r(x) = \text{dist}(x, \partial Q) \leq l_0\} \subset \bigcup_{m=1}^p \prod_m^{l_m, h}$.

Easily verified that for all $x = (x', x_n) \in \Pi_m^{l_m, h}$, $m = 1, \dots, p$,

$$r(x) \leq x_n - \varphi_m(x') \leq \frac{\sqrt{5}}{2} r(x).$$

We fix an index m , $1 \leq m \leq p$, and take a local coordinate system with origin at x^m .

Now define mappings L and L_{-1} of the space R^n onto itself by relations $L(x) = (x', x_n - \varphi_m(x'))$, where $x = (x', x_n)$ and $L_{-1}(y) = (y', y_n + \varphi_m(y'))$ with $y = (y', y_n)$. The image of $\Pi_m^{l_m, h}$ under the mapping L will be denoted by $\tilde{\Pi}_m^{l_m, h}$: $L(\tilde{\Pi}_m^{l_m, h}) = \tilde{\Pi}_m^{l_m, h}$.

Consider the sequence of operators

$$T_k u = (\bar{b}_k(x), \nabla u(x)) - \operatorname{div}(\bar{c}_k(x)u(x)) + d_k(x)u(x), \quad u \in \overset{\circ}{W}_2^1(Q), \quad k = 1, 2, \dots$$

$$\bar{b}_k(x) = \begin{cases} \bar{b}(x), & \text{if } r(x) > \frac{1}{k}, \\ 0, & \text{if } r(x) \leq \frac{1}{k}, \end{cases} \quad \bar{c}_k(x) = \begin{cases} \bar{c}(x), & \text{if } r(x) > \frac{1}{k}, \\ 0, & \text{if } r(x) \leq \frac{1}{k}, \end{cases} \quad d_k(x) = \begin{cases} d(x), & \text{if } r(x) > \frac{1}{k}, \\ 0, & \text{if } r(x) \leq \frac{1}{k}. \end{cases}$$

It can be readily verified that the operator T_k is a compact linear operator from $\overset{\circ}{W}_2^1(Q)$ into $\overset{\circ}{W}_2^{-1}(Q)$. Indeed, let $\{w(x)\}$ be a bounded set in $\overset{\circ}{W}_2^1(Q)$. Then sets $\{(\bar{b}_k(x), \nabla w(x))\}$, $\{\bar{c}_k(x)w(x)\}$ and $\{d_k(x)w(x)\}$ are bounded in $L_2(Q)$ and by that are compact in $\overset{\circ}{W}_2^{-1}(Q)$ (see, for example, [4]). Hence, for the proof of the theorem it is sufficient to show that $\|T - T_k\| \rightarrow 0$ as $k \rightarrow \infty$.

Without loss of generality we suggest that $k > \frac{1}{l_0}$ and functions $\omega_1(t)$, $\omega_2(t)$ are positive. In view of (1) there exists a monotone function $\varepsilon(t) \rightarrow 0$, as $t \rightarrow +0$, such that $|\bar{b}(x)| \leq \frac{\varepsilon(r(x))}{r(x)}$. For $u \in \overset{\circ}{W}_2^1(Q)$ and $\eta \in C_0^\infty(Q)$ consider

$$\langle (T - T_k)u, \eta \rangle = \int_{Q^{j/k}} \left((\bar{b}(x), \nabla u(x))\eta(x) + (\bar{c}(x)u(x), \nabla \eta(x)) + d(x)u(x)\eta(x) \right) dx.$$

Denote $u(y', y_n + \varphi(y')) = \tilde{u}(y)$, $\eta(y', y_n + \varphi(y')) = \tilde{\eta}(y)$.

Due to (1), (2) and (3), we have

$$\begin{aligned} |\langle (T - T_k)u, \eta \rangle| &\leq \int_{Q^{j/k}} \left(\frac{\varepsilon(r(x))|\nabla u(x)||\eta(x)|}{r(x)} + C(r(x))|u(x)||\nabla \eta(x)| + D(r(x))|u(x)||\eta(x)| \right) dx \leq \\ &\leq \varepsilon \left(\frac{1}{k} \right) \int_{Q^{j/k}} \frac{|\nabla u(x)||\eta(x)|}{r(x)} dx + \omega_1^{1/2} \left(\frac{1}{k} \right) \int_{Q^{j/k}} \frac{C(r(x))}{\omega_1^{1/2}(r(x))} |u(x)||\nabla \eta(x)| dx + \\ &\quad + \omega_2^{1/2} \left(\frac{1}{k} \right) \int_{Q^{j/k}} \frac{D(r(x))}{\omega_2^{1/2}(r(x))} |u(x)||\eta(x)| dx. \end{aligned} \tag{4}$$

Let us estimate

$$I_1 = \int_{Q^{1/k}} \frac{|\nabla u(x)| |\eta(x)|}{r(x)} dx \leq \int_{Q^0} \frac{|\nabla u(x)| |\eta(x)|}{r(x)} dx \leq \sum_{m=1}^p \int_{\Pi_m^{l_m, h}} \frac{|\nabla u(x)| |\eta(x)|}{r(x)} dx.$$

In view of the Hardy inequality (see, for example, [5]) for $m=1, \dots, p$ the following estimate holds:

$$\begin{aligned} \int_{\Pi_m^{l_m, h}} \frac{|\nabla u(x)| |\eta(x)|}{r(x)} dx &\leq \sqrt{\frac{5}{2}} \int_{\tilde{\Pi}_m^{l_m, h}} \frac{|\nabla \tilde{u}(y)| |\tilde{\eta}(y)|}{y_n} dy \leq \sqrt{\frac{5}{2}} \left(\int_{\tilde{\Pi}_m^{l_m, h}} |\nabla \tilde{u}(y)|^2 dy \right)^{1/2} \left(\int_{\tilde{\Pi}_m^{l_m, h}} \frac{\tilde{\eta}^2(y)}{y_n^2} dy \right)^{1/2} \leq \\ &\leq \sqrt{5} \left(\int_{\Pi_m^{l_m, h}} |\nabla u(x)|^2 dx \right)^{1/2} \left(\int_{\tilde{\Pi}_m^{l_m, h}} \frac{\tilde{\eta}^2(y)}{y_n^2} dy \right)^{1/2} \leq \text{const} \|u\|_{W_2^1(Q)} \|\eta\|_{W_2^1(Q)}. \end{aligned}$$

Thus,

$$I_1 \leq \text{const} \|u\|_{W_2^1(Q)} \|\eta\|_{W_2^1(Q)}, \quad (5)$$

where the constant does not depend on u and η .

$$\begin{aligned} \text{Next } I_2 &= \int_{Q^{1/k}} \frac{C(r(x))}{\omega_1^2(r(x))} |u(x)| |\nabla \eta(x)| dx \leq \int_{Q^0} \frac{C(r(x))}{\omega_1^{1/2}(r(x))} |u(x)| |\nabla \eta(x)| dx \leq \\ &\leq \sum_{m=1}^p \int_{\Pi_m^{l_m, h}} \frac{C(r(x))}{\omega_1^{1/2}(r(x))} |u(x)| |\nabla \eta(x)| dx. \end{aligned}$$

For $m=1, \dots, p$ we have

$$\begin{aligned} \int_{\Pi_m^{l_m, h}} \frac{C(r(x))}{\omega_1^{1/2}(r(x))} |u(x)| |\nabla \eta(x)| dx &\leq \left(\int_{\Pi_m^{l_m, h}} \frac{C^2(r(x))}{\omega_1(r(x))} u^2(x) dx \right)^{1/2} \|\eta\|_{W_2^1(Q)} \leq \\ &\leq \left(\int_{\tilde{\Pi}_m^{l_m, h}} \frac{C^2\left(\frac{2}{\sqrt{5}}y_n\right)}{\omega_1\left(\frac{2}{\sqrt{5}}y_n\right)} \tilde{u}^2(y) dy \right)^{1/2} \|\eta\|_{W_2^1(Q)} \leq \left(\int_{\tilde{\Pi}_m^{l_m, h}} \frac{C^2\left(\frac{2}{\sqrt{5}}y_n\right)}{\omega_1\left(\frac{2}{\sqrt{5}}y_n\right)} y_n \int_0^{y_n} |\nabla \tilde{u}(y', \tau)|^2 d\tau dy \right)^{1/2} \|\eta\|_{W_2^1(Q)} \leq \\ &\leq \left(\int_0^h dy_n \frac{C^2\left(\frac{2}{\sqrt{5}}y_n\right)}{\omega_1\left(\frac{2}{\sqrt{5}}y_n\right)} y_n \int_{|y'| < l_m} dy' \int_0^h d\tau |\nabla \tilde{u}(y', \tau)|^2 \right)^{1/2} \|\eta\|_{W_2^1(Q)} \leq \\ &\leq \sqrt{2} \left(\int_0^h \frac{C^2\left(\frac{2}{\sqrt{5}}y_n\right)}{\omega_1\left(\frac{2}{\sqrt{5}}y_n\right)} y_n dy_n \right)^{1/2} \|u\|_{W_2^1(Q)} \|\eta\|_{W_2^1(Q)}. \end{aligned}$$

Thus, we get

$$I_2 \leq \text{const} \|u\|_{W_2^1(Q)} \|\eta\|_{W_2^1(Q)}, \quad (6)$$

where the constant does not depend on u and η .

Similarly we obtain

$$I_3 = \int_{Q^{j/k}} \frac{D(r(x))}{\omega_2^{1/2}(r(x))} |u(x)||\eta(x)| dx \leq \int_{Q^0} \frac{D(r(x))}{\omega_2^{1/2}(r(x))} |u(x)||\eta(x)| dx \leq \sum_{m=1}^p \int_{\tilde{\Pi}_m^{j/k}} \frac{D(r(x))}{\omega_2^{1/2}(r(x))} |u(x)||\eta(x)| dx.$$

Finally, for $m = 1, \dots, p$ we get

$$\begin{aligned} \int_{\tilde{\Pi}_m^{j/k}} \frac{D(r(x))}{\omega_2^{1/2}(r(x))} |u(x)||\eta(x)| dx &\leq \int_{\tilde{\Pi}_m^{j/k}} \frac{D\left(\frac{2}{\sqrt{5}} y_n\right)}{\omega_2^{1/2}\left(\frac{2}{\sqrt{5}} y_n\right)} |\tilde{u}(y)||\tilde{\eta}(y)| dy \leq \\ &\leq \left(\int_{\tilde{\Pi}_m^{j/k}} \frac{D^2\left(\frac{2}{\sqrt{5}} y_n\right)}{\omega_2\left(\frac{2}{\sqrt{5}} y_n\right)} y_n^2 \tilde{u}^2(y) dy \right)^{1/2} \left(\int_{\tilde{\Pi}_m^{j/k}} \frac{\tilde{\eta}^2(y)}{y_n^2} dy \right)^{1/2} \leq \\ &\leq \text{const} \left(\int_{\tilde{\Pi}_m^{j/k}} \frac{D^2\left(\frac{2}{\sqrt{5}} y_n\right)}{\omega_2\left(\frac{2}{\sqrt{5}} y_n\right)} y_n^3 \int_0^{y_n} |\nabla \tilde{u}(y', \tau)|^2 d\tau dy \right)^{1/2} \|\eta\|_{W_2^1(Q)} \leq \\ &\leq \text{const} \left(\int_0^h \frac{D^2\left(\frac{2}{\sqrt{5}} y_n\right)}{\omega_2\left(\frac{2}{\sqrt{5}} y_n\right)} y_n^3 dy_n \right)^{1/2} \|u\|_{W_2^1(Q)} \|\eta\|_{W_2^1(Q)}. \end{aligned}$$

Thus,

$$I_3 \leq \text{const} \|u\|_{W_2^1(Q)} \|\eta\|_{W_2^1(Q)}, \tag{7}$$

where the constant does not depend on u and η . From (4)–(7) we obtain the estimate

$$|\langle (T - T_k)u, \eta \rangle| \leq \text{const} \left(\varepsilon \left(\frac{1}{k}\right) + \omega_1^{1/2} \left(\frac{1}{k}\right) + \omega_2^{1/2} \left(\frac{1}{k}\right) \right) \|u\|_{W_2^1(Q)} \|\eta\|_{W_2^1(Q)},$$

where the constant does not depend on u and η .

From this it immediately follows that $\|T - T_k\| \leq \text{const} \left(\varepsilon \left(\frac{1}{k}\right) + \omega_1^{1/2} \left(\frac{1}{k}\right) + \omega_2^{1/2} \left(\frac{1}{k}\right) \right)$

and consequently $\|T - T_k\| \rightarrow 0$ as $k \rightarrow \infty$. The Theorem is proved.

Received 15.11.2010

REFERENCES

1. **Dumanyan V.Zh.** Proceedings of the YSU. Physical and Mathematical Sciences, 2011, № 1, p. 3–6.
2. **Ladyzhenskaya O.A., Uraltseva N.N.** Linear and Quasilinear Elliptic Equations. Academic Press, 1968.
3. **Mikhailov V.P.** Partial Differential Equations. 1st ed. M.: Nauka, 1976; English transl. of 1st ed. M.: Mir, 1978.
4. **Mikhailov V.P., Gushchin A.K.** Advanced Topics of “Equations of Mathematical Physics”. Lecture courses of SEC. M., Steklov Mathematical Institute of RAS, 2007.
5. **Maz'ya V.G.** Sobolev Spaces. L.: Leningrad Univ. Press, 1985 (in Russian); English transl. from the Russian by T.O. Shaposhnikova. Berlin–New York: Springer-Verlag, 1985.

Վ. Մ. Դումանյան
Առաջին կարգի գծային դիֆերենցիալ օպերատորների
կոմպակտության մասին

Հոդվածում ուսումնասիրվում են առաջին կարգի անսահմանափակ գործակիցներով գծային դիֆերենցիալ օպերատորներ: Ապացուցվում է այդ օպերատորների կոմպակտությունը:

В. Ж. Думанян
О компактности линейных дифференциальных
операторов первого порядка

В настоящей работе исследуются линейные дифференциальные операторы первого порядка с неограниченными коэффициентами. Установлена компактность рассматриваемых операторов